HPLC Application

ID No.: 20906



Nicotinic acid / Nicotinamide (10 ng/mL) in Human Plasma by Impact on Gemini 3µm C18 100x4.6mm

Gemini® 3 µm C18 110 Å, LC Column 100 x 4.6 mm, Ea

100 x 4.6 mm ID **Dimensions:** Order No: 00D-4439-E0 **Elution Type:** Gradient

Eluent A: 0.1% formic acid Eluent B: Methanol 100%

Gradient	Step No.	Time (min)	Pct A	Pct B
Profile:	1	0	90	10
	2	2.5	10	90
	3	2.6	90	10
	4	4	90	10






Products used in this application:



Flow Rate: 0.6 mL/min Col. Temp.: ambient

Electrospray Mass Spec (ESMS) @ (ambient) **Detection:**

Detector Info:



ANALYTES:

Nicotinamide

Retention Time: 2.51 min

2 Nicotinic acid

Retention Time: 2.68 min

©2025 Phenomenex Inc. All rights reserved.

For more information contact your Phenomenex Representative at support@phenomenex.com



Phenomenex products are available worldwide. www.phenomenex.com

support@phenomenex.com

Sample Preparation Details

for **HPLC** Application ID No.: **20906**



Nicotinic acid / Nicotinamide (10 ng/mL) in Human Plasma by Impact on Gemini 3µm C18 100x4.6mm

PRODUCT DESCRIPTION:

Impact™ Protein Precipitation, 2mL Square Well Filter Plate, 2/Pk

Order No.: CE0-7565

SOLID PHASE EXTRACTION (SPE) PRODCEDURE:

Note: The solvent volumes shown below are for a Proprietary bed mass.

The solvent volumes will need to be adjusted for a smaller or larger bed mass.

Condition:
Load:
Wash:
Dry:
Elute:
Final Prep and Analysis:
Inject: 2 µL on HPLC Electrospray Mass Spec (ESMS) @ (ambient)

ANALYTES:	Spiked Conc.	Log P	рКа	% Rec	%RSC
	(ng/mL)				(n=0)
 Nicotinamide 	10				
2 Nicotinic acid	10				

This method is designed as a convenient starting point for further investigation and can be tailored to meet your extraction goals. Call your local Phenomenex Representative for assistance in method development and optimization techniques.

©2025 Phenomenex Inc. All rights reserved.

For more information contact your Phenomenex Representative at support@phenomenex.com

