

HPLC Application

ID No.: 24526

Separation of Isobaric cannabinoids AB-PINACA and THC-OH on Luna Omega 3µm Polar C18 100x3.0mm

Column: Luna® Omega 3 µm Polar C18 100 Å, LC Column 100 x 3.0 mm, Ea

Dimensions: 100 x 3 mm ID

Order No: 00D-4760-Y0

Elution Type: Gradient

Eluent A: 0.1% formic acid

Eluent B: 0.1% formic acid/methanol

Gradient Profile:	Step No.	Time (min)	Pct A	Pct B
	1	0	90	10
	2	2	5	95
	3	4	5	95
	4	4.01	90	10
	5	6	90	10


Flow Rate: 0.6 mL/min

Col. Temp.: 25 °C

Detection: Tandem Mass Spec (MS-MS) @ (25 °C)
24526



Products used in this application:



ANALYTES:

1 AB-PINACA

Retention Time: 3.15 min

2 THC-OH

Retention Time: 3.38 min



Sample Preparation Details

for HPLC Application ID No.: 24526

Separation of Isobaric cannabinoids AB-PINACA and THC-OH on Luna Omega 3µm Polar C18 100x3.0mm

PRODUCT DESCRIPTION:

Strata™-X-C 33 µm Polymeric Strong Cation, 30 mg / 3 mL, Tubes , 50/Pk

Order No.: 8B-S029-TBJ

SOLID PHASE EXTRACTION (SPE) PROCEDURE:

Note: The solvent volumes shown below are for a 30 mg bed mass.

The solvent volumes will need to be adjusted for a smaller or larger bed mass.

Condition:

Load:

Wash:

Dry:

3-4 mins at 10-15" of Hg

Elute:

Final Prep and Analysis:

Sample Prep Product: Strata-X-C, 30 mg/3 mL (Part No. 8B-S029-TBJ)

Sample pre-treatment

Inject: 20 µL on HPLC Tandem Mass Spec (MS-MS) @ (25°C)

ANALYTES:	Spiked Conc. (ng/mL)	Log P	pKa	% Rec	%RSC (n=0)
1 AB-PINACA	50				
2 THC-OH	50				

Note: This method is designed as a convenient starting point for further investigation and can be tailored to meet your extraction goals. Call your local Phenomenex Representative for assistance in method development and optimization techniques.

©2024 Phenomenex Inc. All rights reserved.

For more information contact your Phenomenex Representative at support@phenomenex.com



Phenomenex products are available worldwide.

www.phenomenex.com

support@phenomenex.com