
HPLC Application

ID No.: 25531

LC-MS/MS Chromatogram for Amlodipine Enantiomers and Internal Standards



ANALYTES:

- 1 (R)-Amlodipine Retention Time: 2.29 min
- 2 (R)-Amlodipine-d4 Retention Time: 2.3 min
- 3 (S)-Amlodipine Retention Time: 2.93 min
- 4 (S)-Amlodipine-d4 Retention Time: 2.94 min

©2025 Phenomenex Inc. All rights reserved.

Phenomenex products are available worldwide.

www.phenomenex.com

For more information contact your Phenomenex Representative at support@phenomenex.com

ohenomenex

for HPLC Application ID No.: 25531



LC-MS/MS Chromatogram for Amlodipine Enantiomers and Internal Standards

PRODUCT DESCRIPTION:

Strata[™]-X 33 µm Polymeric Reversed Phase, 30 mg / 1 mL, Tubes , 100/Pk

Order No.: 8B-S100-TAK

SOLID PHASE EXTRACTION (SPE) PRODCEDURE:

Note: The solvent volumes shown below are for a bed mass.

The solvent volumes will need to be adjusted for a smaller or larger bed mass.

Condition:

Load:

5 μ L of internal standard working solution was added to 100 μ L spiked human plasma (spiked at 0.050, 0.150, 20.0, 40.0, and 50.0 ng/mL) and mixed, 500 μ L of 0.2% (v/v) Ethanolamine in water was added to sample and vortexed.

Wash:			
Dry:			
Elute:			

Final Prep and Analysis:

Inject: 10 μL on HPLC LC/MS/MS @ 0.0000000000 (ambient)

ANALYTES:	Spiked Conc. (ng/mL)	Log P	рКа	% Rec	%RSC (n=0)
1 (R)-Amlodipine	0				
2 (R)-Amlodipine-d4	0				
3 (S)-Amlodipine	0				
4 (S)-Amlodipine-d4	0				

Note: This method is designed as a convenient starting point for further investigation and can be tailored to meet your extraction goals. Call your local Phenomenex Representative for assistance in method development and optimization techniques.

©2025 Phenomenex Inc. All rights reserved.

Phenomenex products are available worldwide.

www.phenomenex.com

For more information contact your Phenomenex Representative at support@phenomenex.com